



**International Science Group**

**ISG-KONF.COM**

**IV**  
**INTERNATIONAL SCIENTIFIC**  
**AND PRACTICAL CONFERENCE**  
**"DISCUSSION AND DEVELOPMENT OF MODERN**  
**SCIENTIFIC RESEARCH"**

**Helsinki, Finland**  
**October 18 - 21, 2022**

**ISBN 979-8-88796-813-1**

**DOI 10.46299/ISG.2022.2.4**

# **DISCUSSION AND DEVELOPMENT OF MODERN SCIENTIFIC RESEARCH**

Proceedings of the IV International Scientific and Practical Conference

Helsinki, Finland  
October 18 – 21, 2022

**UDC 01.1**

The 4th International scientific and practical conference “Discussion and development of modern scientific research” (October 18-21, 2022) Helsinki, Finland. International Science Group. 2022. 487 p.

**ISBN – 979-8-88796-813-1**

**DOI – 10.46299/ISG.2022.2.4**

## EDITORIAL BOARD

<u>Pluzhnik Elena</u>	Professor of the Department of Criminal Law and Criminology Odessa State University of Internal Affairs Candidate of Law, Associate Professor
<u>Liudmyla Polyvana</u>	Department of Accounting and Auditing Kharkiv National Technical University of Agriculture named after Petr Vasilenko, Ukraine
<u>Mushenyk Iryna</u>	Candidate of Economic Sciences, Associate Professor of Mathematical Disciplines, Informatics and Modeling. Podolsk State Agrarian Technical University
<u>Prudka Liudmyla</u>	Odessa State University of Internal Affairs, Associate Professor of Criminology and Psychology Department
<u>Marchenko Dmytro</u>	PhD, Associate Professor, Lecturer, Deputy Dean on Academic Affairs Faculty of Engineering and Energy
<u>Harchenko Roman</u>	Candidate of Technical Sciences, specialty 05.22.20 - operation and repair of vehicles.
<u>Belei Svitlana</u>	Ph.D., Associate Professor, Department of Economics and Security of Enterprise
<u>Lidiya Parashchuk</u>	PhD in specialty 05.17.11 "Technology of refractory non-metallic materials"
<u>Levon Mariia</u>	Candidate of Medical Sciences, Associate Professor, Scientific direction - morphology of the human digestive system
<u>Hubal Halyna Mykolaiivna</u>	Ph.D. in Physical and Mathematical Sciences, Associate Professor

37.	Шнайдер А.О. ПЕРЕВАГИ СІМЕЙНОЇ МЕДІАЦІЇ НАД СУДОВОЮ ФОРМОЮ ВИРІШЕННЯ СПОРІВ	188
MANAGEMENT, MARKETING		
38.	Khmelnitskiy A. MANAGEMENT ACTIVITY TECHNIQUES FOR CREATING AN ANTI-CRISIS MANAGEMENT SYSTEM OF AN EDUCATIONAL INSTITUTION	193
39.	Voronina A., Hanzhuk S., Hanzhuk K. IMPROVEMENT OF THE PROCESS OF MANAGEMENT OF ENTERPRISE ASSETS	196
40.	Василиха Н.В., Тороній Л.І., Боїшко Б.Ю. УПРАВЛІННЯ ЛЮДСЬКИМИ РЕСУРСАМИ – ОСНОВА КОНКУРЕНТОСПРОМОЖНОСТІ ПІДПРИЄМСТВА	201
41.	Заболотна О.С. СУТНІСТЬ КАДРОВОГО ПОТЕНЦІАЛУ ЯК ІНСТРУМЕНТ ВПЛИВУ НА ЕФЕКТИВНІСТЬ ЙОГО ВИКОРИСТАННЯ	204
42.	Микитенко Н.В. СФЕРА РИТЕЙЛУ В УКРАЇНІ: ВОЄННІ РЕАЛІЇ ТА ПЕРСПЕКТИВИ	207
43.	Рябенська М.О., Цимбалюк К.В. АКТУАЛЬНІ ПРОБЛЕМИ ПІДБОРУ КАДРІВ ЯК ВАЖЛИВОГО ФАКТОРУ УПРАВЛІННЯ ПЕРСОНАЛОМ НА ПІДПРИЄМСТВАХ ГОТЕЛЬНОГО ТА РЕСТОРАННОГО БІЗНЕСУ	211
MEDICINE		
44.	Baieva O., Baiev V. THE ESSENCE AND ORIGIN OF MEDICAL INSURANCE	215
45.	Hnidoi I., Hnida N. THE STATE OF PROOXIDANT-ANTIOXIDANT HOMEOSTASIS IN CHILDREN WITH DIFFERENT LEVELS OF LEAD IN THE BLOOD	222

## **THE STATE OF PROOXIDANT-ANTIOXIDANT HOMEOSTASIS IN CHILDREN WITH DIFFERENT LEVELS OF LEAD IN THE BLOOD**

**Hnidoi Ihor,**  
Ph.D., Associate Professor,  
International Humanitarian University, Odesa

**Hnida Natalia,**  
Intern doctor,  
Odesa Regional Children's Clinical Hospital

Lead is an environmental toxicant that can induce oxidative stress (OS) via reactive oxygen species (ROS) generation, which has been reported as an important mechanism underlying lead toxicity [1]. To install biochemical mechanisms of these disorders, the state of prooxidant-antioxidant homeostasis in children with different levels of lead in the blood was studied.

100 children of a random sample aged from 4 to 15 years were examined (average age 11.5 years). Determination of lead was carried out in heparinized venous blood by the method of atomic absorption spectrometry with electrothermal atomization.

The activity of lipid peroxidation processes (LPO) was estimated by accumulation of primary (diene conjugates - DK) and final (malonic dialdehyde - MDA) products of lipoperoxidation in the blood. The state of the antioxidant system was evaluated by glutathione peroxidase (GPO) activity and by the content of sulfhydryl and thiol-disulfide groups of blood.

To determine the dependence of LPO-AOS indicators on the level of lead in the blood, children were divided into 4 groups: 1) group with lead content up to 69  $\mu\text{g/l}$ ,  $n = 84$ ; 2) group with a marginal level, within 70-99  $\mu\text{g/l}$ ,  $n = 29$ ; 3) group with an elevated level, 100-149  $\mu\text{g/l}$ ,  $n = 11$ ; 4) group with a sharply increased level, more than 150  $\mu\text{g/l}$ ,  $n = 6$ .

The results are in table.1. The lowest values of pro-oxidant indicators were in group of children with the lowest levels of lead in the blood, up to 69  $\mu\text{g/l}$ . With promotion accumulation of lead, DK values increased. Similarly, MDA values also increased. In parallel, the indicators of the thiol-disulfide system of antioxidant protection also increased - obviously, to repair the balance prooxidants and antioxidants.

Table 1.  
Indicators of the LPO-AOS system in children of different groups

Indicators	Groups of children			
	1	2	3	4
DK, $\mu\text{mol/l}$	22.18 $\pm$ 1.50	31.38 $\pm$ 3.29	46.13 $\pm$ 5.99*	55.94 $\pm$ 5.75*
MDA, $\mu\text{mol/l}$	182.11 $\pm$ 5.11	184.61 $\pm$ 12.91	233.07 $\pm$ 17.23*	266.51 $\pm$ 26.63*
GPO, $\mu\text{mol/l/sec}$	8.38 $\pm$ 0.50	8.54 $\pm$ 0.82	8.83 $\pm$ 1.06	7.90 $\pm$ 1.86
SH-groups, $\mu\text{mol/l}$	1.11 $\pm$ 0.08	1.47 $\pm$ 0.24	2.24 $\pm$ 0.25*	1.96 $\pm$ 0.27*
S-S-links, $\mu\text{mol/l}$	0.45 $\pm$ 0.06	0.64 $\pm$ 0.21	1.57 $\pm$ 0.24*	1.13 $\pm$ 0.32*
SH/SS index	2.50	2.29	1.42	1.74

Note. \* - The difference is significant ( $p < 0.05$ ) with the indicators of the first group.

But a compensatory increase in the indicators of AOS was revealed, presumably, is not enough. Because the SH/SS ratio decreased while the LPO indicators increased, and the level of lead increased in the blood.

This assumption was confirmed by antioxidant activity enzyme glutathione-peroxidase, which had no significant changes, despite for activation in the chain of prooxidants.

Thus, when the content of lead in the blood of children was increased, the activity of LPO also was increased. Stabilization of redox balance in children with lead levels above the norm occurred no longer at a minimum level, as in control, but at a higher level. It led to a certain exhaustion antioxidant reserve of the body. This is consistent with scientific data that OS occurs when the generation of ROS exceeds the antioxidant system's ability to defend cells against oxidized molecules [1].

#### References:

1. Lopes, A.C.B.A., Peixe, T.S., Mesas, A.E., Paoliello, M.M.B. (2016). Lead Exposure and Oxidative Stress: A Systematic Review. In: de Voogt, P. (eds) Reviews of Environmental Contamination and Toxicology Volume 236. Reviews of Environmental Contamination and Toxicology, vol 236. Springer, Cham. [https://link.springer.com/chapter/10.1007/978-3-319-20013-2\\_3](https://link.springer.com/chapter/10.1007/978-3-319-20013-2_3)