

Synthesis, Crystal Structure, and Spectral Characteristics of *N*-(*tert*-Butyl)aminomethanesulfonic Acid

R. E. Khoma^{a,b}, V. O. Gel'mbol'dt^c, A. A. Ennan^a, V. N. Baumer^{d,e}, and A. N. Puzan^d

^a Physicochemical Institute for Human and Environmental Protection,
Ministry of Education and Science of the National Academy of Sciences of Ukraine,
ul. Preobrazhenskaya 3, Odessa, 65082 Ukraine
e-mail: rek@onu.edu.ua

^b Mechnikov Odessa National University, Odessa, Ukraine

^c Odessa National Medical University, Odessa, Ukraine

^d Institute of Single Crystals of the National Academy of Sciences of Ukraine, Kharkov, Ukraine

^e Karazin Kharkiv National University, Kharkov, Ukraine

Received April 2, 2015

Abstract—A new method of the synthesis of *N*-(*tert*-butyl)aminomethanesulfonic acid in the system SO₂–(CH₃)₃CNH₂–CH₂O–H₂O was developed. The target compound [(CH₃)₃C]NHCH₂SO₃H was characterized by means of X-ray diffraction analysis, IR spectroscopy, and mass spectrometry.

Keywords: sulfur(IV) oxide, paraformaldehyde, primary alkylamine, condensation

DOI: 10.1134/S1070363215100102

Compounds of zwitterionic structure, in particular, aminoalkanesulfonic acids, their derivatives and salts [1–4] are promising components of buffer solutions [1, 2] widely used in biological research for monitoring the medium pH. Moreover, the compounds are valuable as biologically active species exhibiting various types of pharmacological activity [4, 5].

Earlier we developed an original method of synthesis of aminoalkanesulfonic acids *N*-derivatives by an example of sulfur(IV) oxide reaction with a mixture of monoethanolamine and formaldehyde in aqueous solution that led to the formation of *N*-(hydroxyethyl)-aminomethanesulfonic acid [6] (Scheme 1).

In extension of [6] we attempted the preparation of one of the *N*-derivatives of aminoalkanesulfonic acid with the use in the reaction of *tert*-butylamine. In the present report we describe the synthesis of *N*-(*tert*-butyl)aminomethanesulfonic acid and the study of its structure.


The synthesis was accomplished by the reaction of sulfur(IV) oxide with aqueous solution of a mixture of *tert*-butylamine and formaldehyde; *N*-(*tert*-butyl)aminomethanesulfonic acid was obtained in quantitative

yield. The composition and structure of the target compound was confirmed by mass spectrometry, elemental analysis, IR and NMR spectroscopy, and X-ray diffraction analysis.

IR spectroscopy data of the compound indicated the zwitterionic structure similarly as for the earlier investigated analogs [6, 7].

General view of the molecule of *N*-(*tert*-butyl)aminomethanesulfonic acid is presented in Fig. 1; the values of bond lengths and bond angles are listed in Table 1.

Scheme 1.






Fig. 1. General view of the molecule of *N*-(*tert*-butyl)aminomethanesulfonic acid; thermal ellipsoids at the 50% probability level.




Fig. 2. Crystal packing and hydrogen bonds in the molecule of *N*-(*tert*-butyl)aminomethanesulfonic acid. Hydrogen bonds are depicted by dashed lines.

In the crystal packing of the molecules hydrogen bonds $\text{N}-\text{H}\cdots\text{O}$ exist between the ammonium group of one molecule and the oxygen atom of the sulfo group of the neighboring molecule that is bound with the first one through a spiral axis. Such hydrogen bonds led to the formation of infinite chains of molecules stretched along the crystallographic axis $[010]$ (Fig. 2). There were no hydrogen bonds between the neighboring chains, only shortened interactions $\text{C}-\text{H}\cdots\text{O}$ were observed. Characteristics of the hydrogen bonds are given in Table 2.

Hence, a possibility of the direct synthesis of *N*-derivatives of aminomethanesulfonic acid through the condensation accompanied by oxidation $\text{S(IV)}\rightarrow\text{S(VI)}$

Table 1. Bond lengths and bond angles in the structure of *N*-(*tert*-butyl)aminomethanesulfonic acid

| Bond | $d, \text{\AA}$ | Angle | ω, deg | Angle | ω, deg |
|-------------------------|-----------------|----------------------------------|----------------------|----------------------------------|----------------------|
| S^1-O^2 | 1.421(3) | $\text{O}^2\text{S}^1\text{O}^3$ | 113.98(18) | $\text{N}^1\text{C}^5\text{S}^1$ | 111.4(2) |
| S^1-O^3 | 1.453(3) | $\text{O}^2\text{S}^1\text{O}^1$ | 114.63(19) | $\text{C}^3\text{C}^1\text{N}^1$ | 109.1(3) |
| S^1-O^1 | 1.459(3) | $\text{O}^3\text{S}^1\text{O}^1$ | 111.14(16) | $\text{C}^3\text{C}^1\text{C}^2$ | 113.0(3) |
| S^1-C^5 | 1.761(4) | $\text{O}^2\text{S}^1\text{C}^5$ | 104.93(18) | $\text{N}^1\text{C}^1\text{C}^2$ | 108.0(3) |
| N^1-C^5 | 1.488(5) | $\text{O}^3\text{S}^1\text{C}^5$ | 106.29(18) | $\text{C}^3\text{C}^1\text{C}^4$ | 111.0(4) |
| N^1-C^1 | 1.521(5) | $\text{O}^1\text{S}^1\text{C}^5$ | 104.87(18) | $\text{N}^1\text{C}^1\text{C}^4$ | 105.3(3) |
| C^1-C^3 | 1.513(5) | $\text{C}^5\text{N}^1\text{C}^1$ | 117.3(3) | $\text{C}^2\text{C}^1\text{C}^4$ | 110.2(3) |
| C^1-C^2 | 1.524(5) | | | | |
| C^1-C^4 | 1.532(5) | | | | |

was confirmed by an example of the system $\text{SO}_2-(\text{CH}_3)_3\text{CNH}_2-\text{CH}_2\text{O}-\text{H}_2\text{O}$.

EXPERIMENTAL

Commercially available sulfur(IV) oxide was used after preliminary purification and drying according to the described method [9]. *tert*-Butylamine and paraformaldehyde of "pure" grade were commercially available reagents and were used without additional purification.

IR spectra were recorded on a Spectrum BX II FT-IR System (Perkin-Elmer) spectrophotometer in the range of $4000-350 \text{ cm}^{-1}$ from KBr pellets. Mass spectra were registered on a MX-1321 instrument (direct input of the sample into the ion source, energy of ionizing electrons 70 eV). Elemental analysis on carbon, hydrogen, and nitrogen was done on a CNH-

Table 2. Characteristics of hydrogen bonds $\text{D}-\text{H}\cdots\text{A}$ in the molecule of *N*-(*tert*-butyl)aminomethanesulfonic acid

| $\text{D}-\text{H}\cdots\text{A}$ | Distance, \AA | | | Angle DHA, deg | Coordinates of atom A |
|--|------------------------|-----------------------------|-----------------------------|----------------------|---|
| | $d(\text{D}-\text{H})$ | $d(\text{H}\cdots\text{A})$ | $d(\text{D}\cdots\text{A})$ | | |
| $\text{N}^1\text{H}^{1\text{A}}\cdots\text{O}^1$ | 0.82(3) | 2.03(4) | 2.829(4) | 166(4) | $-x + 1/2,$ $y + 1/2,$ $-z + 1/2$ |
| $\text{N}^1\text{H}^{1\text{B}}\cdots\text{O}^3$ | 0.93(3) | 1.93(3) | 2.862(4) | 175(3) | $-x + 1/2,$ $y - 1/2,$ $-z + 1/2$ |