

Savytskyi I. V., Znamerovsky S. G., Dzygal O. F. The role of lipid peroxidation in the pathogenesis of experimental peritonitis. *Journal of Education, Health and Sport*. 2020;10(7):389-395. eISSN 2391-8306. DOI <http://dx.doi.org/10.12775/JEHS.2020.10.07.041> <https://apcz.umk.pl/czasopisma/index.php/JEHS/article/view/JEHS.2020.10.07.041> <https://zenodo.org/record/4173559>

The journal has had 5 points in Ministry of Science and Higher Education parametric evaluation. § 8.2) and § 12.1.2) 22.02.2019.
© The Authors 2020;

This article is published with open access at Licensee Open Journal Systems of Nicolaus Copernicus University in Torun, Poland
Open Access. This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. This is an open access article licensed under the terms of the Creative Commons Attribution Non commercial license Share alike. (<http://creativecommons.org/licenses/by-nc-sa/4.0/>) which permits unrestricted, non commercial use, distribution and reproduction in any medium, provided the work is properly cited.
The authors declare that there is no conflict of interests regarding the publication of this paper.

Received: 19.06.2020. Revised: 09.07.2020. Accepted: 31.07.2020.

UDK 616-092:616-036.8

THE ROLE OF LIPID PEROXIDATION IN THE PATHOGENESIS OF EXPERIMENTAL PERITONITIS

I. V. Savytskyi, S. G. Znamerovsky, O. F. Dzygal

Odessa National Medical University, Odessa, Ukraine

Abstract

In the first hour simulated fecal peritonitis development in the rats blood revealed an increase in the level of diene conjugates and malondialdehyde ($p < 0.001$ compared with the results of intact animals. In second stage was detected more pronounced increase in the level of primary and secondary products of lipid peroxidation ($p < 0.001$) as compared with the data of the same hole in the first stage, and compared with the results of group No. 1). On the 3rd day of simulated peritonitis development in rats there is a marked deterioration of the animal condition, which was manifested in lipid peroxidation progression: the diene conjugates level increased at the level of significance $p < 0,001$, and the level of malondialdehyde compared to $p < 0,05$ data from the previous stage of the same group.

Keywords: experimental peritonitis; lipid peroxidation; malonic dialdehyde; diene conjugates.

РОЛЬ ПЕРЕКИСНОГО ОКИСЛЕННЯ ЛІПІДІВ У ПАТОГЕНЕЗІ ЕКСПЕРИМЕНТАЛЬНОГО ПЕРИТОНІТУ

І. В. Савицький, С. Г. Знамеровський, О. Ф. Дзигал

Резюме

На 1-у годину розвитку змодельованого калового перитоніту у крові щурів виявлене збільшення рівня дієнових кон'югатів та малонового діальдегіду ($p < 0,001$ у порівнянні з результатами інтактних тварин). На 2-у етапі виявлене більш виражене підвищення рівня первинних та вторинних продуктів перекисного окислення ліпідів ($p < 0,001$ як у порівнянні з даними цієї ж групи на першому етапі, так і порівняно з результатами групи №1). На 3-ю добу розвитку змодельованого перитоніту у щурів спостерігається виражене погіршення стану тварин, яке проявилось у прогресуванні перекисного окислення ліпідів: рівень дієнових кон'югатів підвищений на рівні значущості $p < 0,001$, а рівень малонового діальдегіду – на рівні $p < 0,05$ у порівнянні з даними попереднього етапу цієї ж групи.

Ключові слова: експериментальний перитоніт; перекисне окислення ліпідів; малоновий діальдегід; дієнові кон'югати.

Introduction

Peritonitis is one of the most severe complications of abdominal pathology [1, 2], which is characterized by a high mortality rate [3, 4].

Violation of the natural mechanisms of detoxification (immune, detoxification functions of the liver and kidneys), which occur in acute peritonitis, due to primary damage to their structure and the development of maladaptive mechanisms, is accompanied by a syndrome of endogenous intoxication [5, 6] This syndrome is a complex of pathological conditions of different etiology and severity of the course, but due to the accumulation of endotoxins in biological fluids and tissues [7]. One of the characteristic and aggravating manifestations of endogenous intoxication syndrome is the activation of lipid peroxidation (LPO) [8]. LPO is one of the key mechanisms for the formation of toxic substances. Having a significant reactivity, the products of lipid peroxidation interact with nucleic acids, proteins and other cell molecules. This leads to uncoupling of oxidative phosphorylation, inactivation of enzymes, and an increase in the permeability of cell membranes [9]. It is also known that oxidized forms of lipids, which accumulate in the blood, lead to destructive processes at the

molecular (free-radical modification of blood lipoproteins), cellular (changes in the functions and structures of biomembranes) and organ (violation of homeostasis) levels [10].

Considering the above, it is relevant to study LPO against the background of the development of peritonitis for the further development of new effective methods for correcting the pathogenetic links of this pathology.

Purpose: to study the dynamics of primary and secondary products of lipid peroxidation in experimental peritonitis.

Materials and methods

The study was performed on 140 white male rats weighing 180–220 g. The animals were divided into 2 groups:

1 group - 40 intact animals

2 group - 100 rats with simulated fecal peritonitis.

Fecal peritonitis was simulated by introducing a 10% fecal suspension at a dose of 0.5 ml per 100 g of animal weight to the abdominal cavity by the puncture method (Lazarenko VA et al., 2016, patent № 233826).

The analysis of indicators was performed for the first hour after the completion of simulation of peritonitis and the first, third and seventh days of the experiment. The research was conducted in accordance with the "Rules of performance of works using experimental animals", approved by the Order of the Ministry of Health of Ukraine № 249 from 01.03.2012 and the Law of Ukraine № 3447-IV "On protection of animals from cruel treatment" (as amended from 15.12.2009 and from 16.10.2012).

Serum levels of diene conjugates (DCs) and malonic dialdehyde (MDA) were determined spectrophotometrically. MDA was determined by reaction with thiobarbituric acid.

Before using parametric methods based on the normality of the statistical distribution, the investigated series of quantitative data were tested for normality using the Shapiro-Wilk's W test. Due to the normal distribution of digital data, Student's parametric criterion was used in the samples.

Research results and their discussion

In the group of animals that simulated fecal peritonitis (taking into account the withdrawal from the experiment of 20 animals at each stage), the following dynamics of survival was observed: 1st hour - 98 rats; 1st day - 49 rats; 3rd day - 20 rats; Day 7 - rats with simulated fecal peritonitis without correction did not survive to this stage of the study.

The choice of DC and MDA as markers of the severity of the pathological process is justified by the following. Data on the content of LPS products have diagnostic value in terms of depth, severity and prognosis of the pathological process. In the development of many diseases, lipoperoxidation can be considered as a universal non-specific pathogenetic link [11]. Lipid peroxidation under physiological conditions regulates the functions of organs and cells. A necessary condition for this is a balance between free radicals and the state of the antioxidant system. That is why the activation of lipid peroxidation plays a negative role in the transition from physiological to pathological, while disrupting regulatory links. Lipid peroxidation itself goes through several stages. It should be noted that each of these stages in endotoxemia may play different roles in the violation of homeostasis. For example, the action of diene conjugates is short-lived, and Schiff alkalis, which act as markers of LPS activity, are chemically inert [12].

In the analysis of diene conjugates, which are the primary products of lipid peroxidation, the following results were obtained in the course of our study (Table 1).

Table 1. The results of the study of the level of diene conjugates in the blood of rats with simulated fecal peritonitis and in intact animals ($M \pm m$)

	1 group	2 group
1st hour	50,41±1,67	68,34±1,98
1st day	50,4±1,98	92,1±1,62
3rd day	50,4±2,27	104,1±1,55
7th day	50,41±1,64	-

In the first stage (the first hour after the completion of peritonitis modeling), an increase in the level of diene conjugates ($p < 0.001$) was found in the blood of rats compared with the results of intact animals.

In the second stage (the first day of development of simulated peritonitis) revealed a more pronounced increase in the level of primary products of lipid peroxidation ($p < 0.001$ both compared with the data of the same group in the first stage and compared with the results of group №1).

On the third day of development of simulated fecal peritonitis in the group of rats without correction there is a marked deterioration of the animals, which manifested itself in the progression of lipid peroxidation: the level of diene conjugates increased at the level of significance $p < 0.001$.

Next, consider the dynamics of the level of malonic dialdehyde (secondary product of LPO) against the background of the development of fecal peritonitis (Table 2).

Table 2. The results of the study of the content of malonic dialdehyde in the blood of rats with simulated fecal peritonitis and in intact animals ($M \pm m$)

	1 group	2 group
1st hour	5,01±0,67	9,24±0,62
1st day	5,01±0,64	16,82±0,49
3rd day	5,01±0,78	19,11±0,59
7th day	5,01±0,67	-

Malonic dialdehyde has characteristic toxic properties, is able to damage nucleic acids, proteins and enzymes and is an indicator of the activity of free radical reactions [12]. Therefore, we can say that excessive accumulation of MDA significantly enhances endotoxemia [13].

During the study of patients with progressive pancreatitis, acute peritonitis, sepsis, ie in critical conditions, an increase in the level of MDA in the blood as the severity of multiorgan failure increased. In the most seriously ill patients, a significant decrease in the levels of antioxidant and proteolytic activities was observed against the background of the maximum increase in the level of MDA [14].

At 1 hour after the completion of modeling peritonitis, an increase in the level of malonic dialdehyde ($p < 0.001$) was found in the blood of rats compared with the results of intact animals.

In the second stage (the first day of development of simulated peritonitis) revealed a more pronounced increase in the level of secondary products of lipid peroxidation ($p < 0.001$ both compared with the data of the same group in the first stage and compared with the results of group №1).

On the third day of development of simulated fecal peritonitis in the group of rats without correction there is a marked deterioration in animals, which manifested itself in the progression of lipid peroxidation: the level of malonic dialdehyde is increased at a level of $p < 0.05$ compared with the same group in the previous stage.

The earliest biochemical indicator, which indicates the beginning of the destructive process in the abdominal cavity, and subsequently reflects the dynamics of the pathological process, is an increase in the concentration of LPO products in the blood [15]. The above

results of our study indicate a pronounced activation of lipid peroxidation on the background of simulated fecal peritonitis in experimental animals.

Conclusions:

- As a result of our study, a pronounced activation of lipid peroxidation on the background of simulated fecal peritonitis in experimental animals.
- Already in the first hour of development of the simulated experimental peritonitis in the blood of rats revealed an increase in the level of diene conjugates and malonic dialdehyde ($p < 0.001$) compared with the results of intact animals)
- The second stage (the first day of development of simulated peritonitis) revealed a more pronounced increase in the level of primary and secondary products of lipid peroxidation ($p < 0.001$ both compared with the data of the same group in the first stage and compared with the results of group №1).
- On the third day of development of simulated fecal peritonitis in the group of rats without correction there is a marked deterioration in animals, which manifested itself in the progression of lipid peroxidation: the level of diene conjugates increased at the level of $p < 0.001$, and the level of malonic dialdehyde at the level of $p < 0.05$ in comparison with the data of the previous stage of the same group.

List of literature sources:

1. Polyansky IY Treatment of acute peritonitis: the transition from evidence to personalized medicine on the basis of genetic research / IY Polyansky, VI Moskalyuk, PV Moroz [etc.]// Art of medicine. – 2018. – №4(8). – P. 148-151. (in Ukrainian)
2. Fofanov OD Treatment of meconium peritonitis: problems and ways to solve them / OD Fofanov, VO Fofanov, RI Nikiforuk // Pediatric surgery. – 2017. – №3 (56). – P. 61-67. (in Ukrainian)
3. Dzyubanovsky I. Ya. Peculiarities of the state of the prooxidant system in experimental peritonitis on the background of diabetes / I. Ya. Dzyubanovsky, BM Vervega, SR Podruchna [etc.] // Medical and clinical chemistry. – 2018. – T. 20. – № 4. – P. 66-71. (in Ukrainian)
4. Fomin PD Tertiary peritonitis as a problem of abdominal surgery / PD Fomin, OB Matviychuk // Clinical surgery. – 2018. - T. 85, № 1. - P. 49-51. (in Ukrainian)
5. Popov AN Endotoxemia in experimental acholia. / A. N. Popov, M. M. Minnebaev. // Bulletin of Experimental Biology and Medicine. – 1997. – № 1. – P. 101 - 102.

(in Russian)

6. Eryukhin I. A. Surgical sepsis (debatable aspects of the problem) / I. A. Eryukhin, S.A. Shlyapnikov // *Surgery*. – 2000. – № 3. – P. 44-46. (in Russian)

7. Morris I. T., Kelty W. *South med. J.* - 1992. - Vol. 85, N 10. - P. 1030-1031.

8. Plum J. Nitric oxide production in peritoneal macrophages from peritoneal dialysis patients with bacterial peritonitis / J. Plum, M.M. Tabatabaei, M.R. Lordnejad [et al.] // *Perit-Dial-Int.* – 1999; 19 Suppl 2. – P.378-83.

9. Eryukhin I. A. Inflammation as a general clinical reaction (based on the model of acute peritonitis) / I. A. Eryukhin, V. Ya. Bely, V. K. Wagner // *St. Petersburg: Science*, 1989. – 262 p. (in Russian)

10. Vlasov AP Functional state of platelets and lipid peroxidation of blood in patients with acute destructive pancreatitis / AP Vlasov, RS Ashirov, VA Trofimov [and others] // *Reanimatology at the turn of the XXI century. (Materials of the International Symposium dedicated to the 60th anniversary of the Research Institute of General Reanimatology of the Russian Academy of Medical Sciences.)* December 2-4, 1996. Moscow. – P. 259-261. (in Russian)

11. Zhdanov GG The problem of hypoxia in intensive care patients in the light of free radical theory / GG Zhdanov, ML Nodel. // *Bulletin of intensive care.* - 1996. – № 1. – P. 23-27. (in Russian)

12. Konyukhova S. G. Effect of hemosorption on the activity of free radical oxidation in destructive diseases of the abdominal organs / S. G. Konyukhova, A. Yu. Dubikaitis, V. I. Strashnov [and others] // *Anesthesiology and resuscitation.* – 1990. – № 6. – P. 48-52. (in Russian)

13. Marusanov VE The role of changes in the blood of biologically active substances, LPO products, thiol-disulfide and ascorbate metabolism in the formation of the severity of endogenous intoxication / VE Marusanov, A.B. Bichun, A.D. Bulganin // *Endogenous intoxication. Abstracts of the International Symposium June 14-16 1994 y.-S.-P.-1994.* – p. 39. (in Russian)

14. Konyukhova S. G. Influence of hemosorption on the activity of free radical oxidation in destructive diseases of the abdominal organs / S. G. Konyukhova, A. Yu. Dubikaitis, V. I. Strashnov [and others] // *Anesthesiology and resuscitation.* – 1990. – № 6. – P.48-52. (in Russian)